

UNCLASSIFIED

AD NUMBER
AD487274
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; Jul 1966. Other requests shall be referred to Army Biological Defense Research Center, Frederick, MD.
AUTHORITY
BDRL D/A ltr, 28 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

487274
TECHNICAL MANUSCRIPT 311

GENE TRANSFER BY CONJUGATION
BETWEEN P. PSEUDOTUBERCULOSIS F-LAC
AND P. PESTIS

William D. Lawton
Brian C. Morris
Thomas W. Burrows

July 1966

UNITED STATES ARMY
BIOLOGICAL CENTER
FORT DETRICK

Reproduction of this publication in whole or part is prohibited except with permission of Commanding Officer, U. S. Army Biological Center, ATTN: Technical Releases Group, Technical Information Department, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

BLANK PAGE

U.S. ARMY BIOLOGICAL CENTER
Fort Detrick, Frederick, Maryland

TECHNICAL MANUSCRIPT 311

GENE TRANSFER BY CONJUGATION BETWEEN
P. PSEUDOTUBERCULOSIS F-LAC AND P. PESTIS

William D. Lawton

Brian C. Morris

Thomas W. Burrows

Medical Bacteriology Department
BIOLOGICAL SCIENCES LABORATORY

Project 1C522301A059

July 1966

FOREWORD

This work was performed during a one-year visit of W.D. Lawton to Microbiological Research Establishment, Porton, Salisbury, Wilts, United Kingdom. The junior authors are on the staff of M.R.E.

ACKNOWLEDGMENT

This investigation was aided by the excellent technical assistance of P.E. Stone.

ABSTRACT

Introduction of the F-lac episome from Escherichia coli to Pasteurella pseudotuberculosis yielded a strain of P. pseudotuberculosis F-lac that could transfer its genes by conjugation to Pasteurella pestis. The selected P. pseudotuberculosis marker arginine⁺ was transferred at frequencies between 10^{-6} and 10^{-7} per donor cell. Unselected donor characters such as melibiose or rhamnose fermentation, phage sensitivity, and urease production appeared at different frequencies in the recombinants. The transferred markers were unrelated to the transfer of F-lac or its subsequent loss from the recipient cell.

I. INTRODUCTION

Introduction of the F-lac episome from Escherichia coli into Pasteurella pseudotuberculosis established a gene transfer system between P. pseudotuberculosis F-lac and various auxotrophs of P. pseudotuberculosis.¹ Since P. pseudotuberculosis is closely related to Pasteurella pestis, we investigated the possibility of extending the F-lac - mediated gene transfer system to include P. pestis.

II. MATERIALS AND METHODS

A. BACTERIAL STRAINS

The donor strain of P. pseudotuberculosis 32/IV F-lac cys⁻, pur⁻, thi⁻ was described previously.¹ The recipient strain of P. pestis was obtained as follows: (i) a methionine-, valine-, isoleucine-independent isolate² was selected from the virulent Alexander strain; (ii) after treatment with ultraviolet irradiation, an arginine E⁻ mutant was obtained; (iii) an isolate resistant to 1000 µg of streptomycin sulfate per ml was picked; (iv) a VW antigen³ calcium requirement⁻³ avirulent colony was isolated. The final strain, labeled P. pestis 370, required cysteine (or thiosulfate), phenylalanine, and arginine for growth, was streptomycin-resistant and pigmented, produced fraction I antigen and pesticin I, and had the same gross antigens, phage sensitivity, and carbohydrate fermentation reactions as a classical strain of P. pestis.

B. MEDIA

Tryptic digest meat broth (TMB) or agar (TMA), Difco heart infusion broth (HIB), and Difco blood agar base (BAB) were employed as complete media. The basic minimal medium employed¹ was supplemented with the following (final concentrations expressed in millimoles in parentheses): cysteine (0.4); phenylalanine (0.5); methionine (0.2); valine (0.2); isoleucine (0.5); glycine (1.0); and sodium metabisulfite (0.5).

C. ABBREVIATIONS

The following abbreviations are used in this report: lac, lactose; arg, arginine; cys, cysteine; phe, phenylalanine; met, methionine; val, valine; ile, isoleucine; gly, glycine; mel, melibiose; rha, rhamnose; ure, urease; oIV³, sensitivity to a bacteriophage originating as a

host-range mutant of a phage culture on P. pestis, serially propagated on P. pseudotuberculosis strain 32/IV; T7^r, resistance to phage T7; Sm^s, sensitivity to 1000 µg of streptomycin sulfate per ml; pig^{Pstb}, forming a large light-brown colony on hemin agar⁴ characteristic of P. pseudotuberculosis in contrast to the compact, dark-brown colonies formed by P. pestis; pes I, pesticin I;⁵ fib, fibrinolysin.⁶

D. MATING PROCEDURE

Essentially, the recipient strain was spread on a minimal agar medium selective for arg⁺ recombinants, and, after ultraviolet irradiation, the donor strain was spread over the recipient lawn. Details of the mating procedure are described in previous publication.¹

III. RESULTS

A. GENE TRANSFER

The frequency of arg⁺ transfer from P. pseudotuberculosis F-lac to P. pestis 370 ranged from 10^{-6} to 10^{-7} per donor cell, which was about two logs less than we had observed for the transfer of four different markers between P. pseudotuberculosis strains.¹ This low frequency, resulting in only 5 to 50 recombinant colonies on the selective plates, was workable because of the complete absence of donor or recipient reversions on control plates.

Selection for cys⁺ or phe⁺ recombinants yielded no colonies.

B. EFFECT OF MEDIUM ON MATING

Although the recipient strain was cys⁻ phe⁻ arg⁻, no arg⁺ recombinants appeared in a minimal agar medium containing only cys and phe. The addition of met, val, ile, and gly (the typical growth requirements of P. pestis) apparently was necessary for mating. The arg⁺ recombinants grew well on minimal agar containing only cys and phe.

As we had observed with the P. pseudotuberculosis X P. pseudotuberculosis mating system,¹ no recombinants resulted from broth matings.

C. ANALYSIS OF UNSELECTED MARKERS

Although *P. pseudotuberculosis* and *P. pestis* are similar, several easily measured properties can be used to distinguish the two species. Our donor strain of *P. pseudotuberculosis* was *arg*⁺, *mel*⁺, *phiV*^s, *rha*⁺, *ure*⁺, T7E, *ntcPstb*, *Sm*^s, *pes* I⁻, *fib*⁻, *phc*⁺; the recipient strain of *P. pestis* was the opposite for these characters. Analysis of 118 recombinants, obtained from six separate experiments, revealed that *mel*⁺ and *phiV*^s were transferred at frequencies of 39% and 5%, *rha*⁺ and *ure*⁺ were transferred at a frequency of approx 1%, and none of the other donor markers were detected in the recombinants (Table 1).

TABLE 1. ANALYSIS OF *arg*⁺ RECOMBINANTS FOR UNSELECTED MARKERS

<i>P. pseudotuberculosis</i> Donor Characters					Analysis of 118 <i>P. pestis</i> Recombinants ^a	
<i>arg</i> ⁺	<i>mel</i> ⁺	<i>phiV</i> ^s	<i>rha</i> ⁺	<i>ure</i> ⁺	No.	%
+	-	-	-	-	62	52.5
+	+	-	-	-	46	39.0
+	+	+	-	-	6	5.1
+	+	+	+	-	1	0.85
+	+	-	+	-	1	0.85
+	-	-	+	-	1	0.85
+	+	-	-	+	1	0.85

a. Obtained from six separate experiments.

D. F-LAC TRANSFER

Twelve of the 118 recombinants were *lac*⁺, segregating *lac*⁻ clones at different frequencies. The *lac*⁺ marker was distributed at random among the recombinant classes and loss of the *lac*⁺ marker did not change any other character.

E. BASE COMPOSITION

The *T_m* of DNA⁷ extracted from the donor strain of *P. pseudotuberculosis* was 88.5 C (46.8% G + C). The *T_m* of *P. pestis* 370 measured at the same time was 88.4 C (46.6% G + C).

IV. DISCUSSION

The data presented indicate that the episome F-lac can promote chromosome transfer between P. pseudotuberculosis and P. pestis similar to that mediated by F-lac in E. coli.⁶ The frequency of transfer observed so far is low (10^{-6} per donor cell) and only arg⁺ has been used successfully as a selected marker. The detection of four unselected donor characters in the arg⁺ recombinants in decreasing frequencies can be interpreted in terms of classical chromosome transfer.

The conversion of P. pestis to P. pseudotuberculosis has been reported by several Soviet investigators (reviewed by Pollitzer⁸), but never with genetically marked strains, and the experiments described have not dispelled all doubt. If this conversion occurs, it might be expected that the distinguishing properties between the two species would be controlled by a single gene locus. Successful gene transfer of one of these properties, e.g. mel⁺, should be correlated with the transfer of other properties such as rha⁺ and ure⁺. The fact that these properties are not jointly transferred (Table 1) suggests that the main properties distinguishing P. pestis and P. pseudotuberculosis are not controlled by the same gene locus.

Investigation of the genetics of virulence requires a reproducible disease syndrome plus a gene transfer system. Plague can be considered a "model" acute disease because the human symptoms occur in common laboratory animals after injection of a few microorganisms. The etiological agent, P. pestis, has been the subject of extensive research and a great deal is known about virulence factors and immunity in this pathogen.⁹⁻¹¹ The extension of a gene transfer system to include P. pestis should lead to a resolution of some of the problems concerning virulence in this pathogen and eventually contribute toward a better understanding of virulence at the genetic level.

LITERATURE CITED

1. Lawton, William D.; Morris, Brian C.; Burrows, Thomas W. October 1965. Gene transfer by conjugation in Pasteurella, (Technical Manuscript 253). Medical Bacteriology Division, U.S. Army Biological Laboratories, Frederick, Maryland.
2. Englesberg, E.; Ingraham, L. 1957. Meiotrophic mutants of Pasteurella pestis and their use in the elucidation of nutritional requirements. Proc. Nat. Acad. Sci. U.S. 43:369-372.
3. Higuchi, K.; Smith, J.L. 1961. Studies on the nutrition and physiology of Pasteurella pestis: VI. A differential plating medium for the estimation of the mutation rate to avirulence. J. Bacteriol. 81:605-608.
4. Jackson, S.; Burrows, T.W. 1956. The pigmentation of Pasteurella pestis on a defined medium containing haemin. Brit. J. Exp. Pathol. 37:570-576.
5. Ben-Gurion, R.; Hertman, I. 1958. Bacteriocin-like material produced by Pasteurella pestis. J. Gen. Microbiol. 19:289-297.
6. Beesley, E.D.; Janssen, W.A.; Surgalla, M.J. 1963. Fibrinolytic activity of Pasteurella pestis. Bacteriol. Proc. p. 80.
7. Marmur, J.; Doty, P. 1962. Determination of the base composition of deoxyribonucleic acid from its thermal denaturation temperature. J. Mol. Biol. 5:109-118.
8. Jacob, F.; Adelberg, E.A. 1959. Transfer of genetic characters by incorporation in the sex factor of Escherichia coli. Compt. Rend. 249:189.
9. Pollitzer, R. 1954. Plague. WHO Monogr. Ser. 22, 698 p.
10. Surgalla, M.J. 1960. Properties of virulent and avirulent strains of Pasteurella pestis. Ann. N.Y. Acad. Sci. 88:1136-1145.
11. Burrows, T.W. 1963. Virulence of Pasteurella pestis and immunity to plague. Ergeb. Mikrobiol. Immunitätsforsch. Exp. Therap. 37:59-113.

Unclassified
Security Classification

DOCUMENT CONTROL DATA - RAD		
(Source: classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1 ORIGINATING ACTIVITY (Corporate author) U.S. Army Biological Center Fort Detrick, Frederick, Maryland, 21701		2a REPORT SECURITY CLASSIFICATION Unclassified 2f GROUP
3 REPORT TITLE GENE TRANSFER BY CONJUGATION BETWEEN <u>P. PSEUDOTUBERCULOSIS F-LAC</u> AND <u>P. PESTIS</u>		
4 DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5 AUTHOR(S) (Last name, first name, initial) Lawton, William D. Morris, Brian C. Burrows, Thomas W.		
6 REPORT DATE July 1966	7a TOTAL NO OF PAGES 10	7b NO OF PAGES 11
8a CONTRACT OR GRANT NO b PROJECT NO 1C522301A059		9a ORIGINATOR'S REPORT NUMBER(S) Technical Manuscript 311 9b OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
10 AVAILABILITY LIMITATION NOTICES Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11 SUPPLEMENTARY NOTES		12 SPONSORING MILITARY ACTIVITY U.S. Army Biological Center Fort Detrick, Frederick, Maryland, 21701
13 ABSTRACT Introduction of the F-lac episome from <u>Escherichia coli</u> into <u>Pasteurella pseudotuberculosis</u> yielded a strain of <u>P. pseudotuberculosis F-lac</u> that could transfer its genes by conjugation to <u>Pasteurella pestis</u> . The selected <u>P. pseudotuberculosis</u> marker arginine ⁺ was transferred at frequencies between 10 ⁻⁶ and 10 ⁻⁷ per donor cell. Unselected donor characters such as melibiose or rhamnose fermentation, phage sensitivity, and urease production appeared at different frequencies in the recombinants. The transferred markers were unrelated to the transfer of F-lac or its subsequent loss from the recipient cell.		

DD FORM 1473
1 JAN 64

Unclassified
Security Classification